

## RESEARCH PAPER

# Programmed cell death activated by Rose Bengal in *Arabidopsis thaliana* cell suspension cultures requires functional chloroplasts

Jorge Gutiérrez<sup>1,2,\*</sup>, Sergio González-Pérez<sup>1,\*</sup>, Francisco García-García<sup>3</sup>, Cara T. Daly<sup>4,5</sup>, Óscar Lorenzo<sup>6</sup>, José L. Revuelta<sup>7</sup>, Paul F. McCabe<sup>5</sup> and Juan B. Arellano<sup>1,†</sup>

<sup>1</sup> Instituto de Recursos Naturales y Agrobiología de Salamanca (IRNASA-CSIC), Cordel de merinas 52, 37008 Salamanca, Spain

<sup>2</sup> Department of Nutritional Sciences, School of Biosciences and Medicine, Faculty of Health and Medical Sciences, University of Surrey, Guildford, Surrey GU2 7XH, UK

<sup>3</sup> Functional Genomics Node, INB, Computational Medicine, Prince Felipe Research Centre, Av. Autopista del Saler 16, Camino de las Moreras, 46012 Valencia, Spain

<sup>4</sup> School of Science, Department of Chemical and Life Sciences, Waterford Institute of Technology, Cork Road, Waterford, Ireland

<sup>5</sup> School of Biology and Environmental Science, University College Dublin, Belfield, Dublin 4, Ireland

<sup>6</sup> Departamento de Fisiología Vegetal, Centro Hispano-Luso de Investigaciones Agrarias, Facultad de Biología, Universidad de Salamanca, C/ Río Duero 12, 37185 Salamanca, Spain

<sup>7</sup> Departamento de Microbiología y Genética, Universidad de Salamanca, Campus Miguel de Unamuno, E-37007 Salamanca, Spain

\* These authors contributed equally to this work.

† To whom correspondence should be addressed. E-mail: [juan.arellano@irnsa.csic.es](mailto:juan.arellano@irnsa.csic.es)

Received 24 January 2014; Revised 25 February 2014; Accepted 12 March 2014

## Abstract

Light-grown *Arabidopsis thaliana* cell suspension culture (ACSC) were subjected to mild photooxidative damage with Rose Bengal (RB) with the aim of gaining a better understanding of singlet oxygen-mediated defence responses in plants. Additionally, ACSC were treated with H<sub>2</sub>O<sub>2</sub> at concentrations that induced comparable levels of protein oxidation damage. Under low to medium light conditions, both RB and H<sub>2</sub>O<sub>2</sub> treatments activated transcriptional defence responses and inhibited photosynthetic activity, but they differed in that programmed cell death (PCD) was only observed in cells treated with RB. When dark-grown ACSC were subjected to RB in the light, PCD was suppressed, indicating that the singlet oxygen-mediated signalling pathway in ACSC requires functional chloroplasts. Analysis of up-regulated transcripts in light-grown ACSC, treated with RB in the light, showed that both singlet oxygen-responsive transcripts and transcripts with a key role in hormone-activated PCD (i.e. ethylene and jasmonic acid) were present. A co-regulation analysis proved that ACSC treated with RB exhibited higher correlation with the conditional fluorescence (*flu*) mutant than with other singlet oxygen-producing mutants or wild-type plants subjected to high light. However, there was no evidence for the up-regulation of *EDS1*, suggesting that activation of PCD was not associated with the EXECUTER- and *EDS1*-dependent signalling pathway described in the *flu* mutant. Indigo Carmine and Methylene Violet, two photosensitizers unable to enter chloroplasts, did not activate transcriptional defence responses in ACSC; however, whether this was due to their location or to their inherently low singlet oxygen quantum efficiencies was not determined.

**Key words:** *Arabidopsis* cell cultures, photosensitizers, programmed cell death, reactive oxygen species, Rose Bengal, singlet oxygen, transcriptional defence responses.

Abbreviations: ACSC, *Arabidopsis thaliana* cell suspension cultures; Chl, chlorophyll; DAB, 3,3'-diaminobenzidine tetrahydrochloride; DNPH, 2,4-dinitrophenylhydrazine; ET, ethylene; EX, EXECUTER; FDA, fluorescein diacetate; HL, high light; IC, indigo carmine; JA, jasmonic acid; LL, low light; MV, Methylene Violet; OPDA, 12-oxophytodienoic acid; PCD, programmed cell death; Pchl, protochlorophyllide; RB, Rose Bengal; ROS, reactive oxygen species; SA, salicylic acid.

© The Author 2014. Published by Oxford University Press on behalf of the Society for Experimental Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

## Introduction

Singlet oxygen ( $^1\text{O}_2$ ) is a reactive oxygen species (ROS) that is formed constitutively in photosystem II (PSII) of plant chloroplasts. Plants can cope with the basal production of  $^1\text{O}_2$  under normal environmental conditions, but high levels of  $^1\text{O}_2$  are produced in response to excess excitation energy in PSII when photosynthetic activity is inhibited by stress or inhibitors (Mullineaux and Baker, 2010). Overproduction of  $^1\text{O}_2$  causes damage to lipids and proteins in the neighbourhood of PSII, leading to a decrease in photosynthetic efficiency and an inhibition of plant growth. Damage to the  $\beta$ -carotene molecules of the PSII reaction centre (RC) by  $^1\text{O}_2$  has also been reported and, interestingly, some of the  $\beta$ -carotene oxidation products have been proposed to be stress signals that mediate gene responses to  $^1\text{O}_2$  (Ramel *et al.*, 2012a, b). The dual role of  $^1\text{O}_2$  as a cytotoxic molecule and a signal molecule is now well established (Kim *et al.*, 2008; Triantaphylidès and Havaux, 2009; Fischer *et al.*, 2013) and much of what is known about the role of  $^1\text{O}_2$  as a signalling molecule comes from the conditional *fluorescence* (*flu*) mutant of *Arabidopsis* (op den Camp *et al.*, 2003; Danon *et al.*, 2005; Laloi *et al.*, 2007; Lee *et al.*, 2007). The *flu* mutant contains a mutation in a negative regulator of chlorophyll (Chl) biosynthesis that results in enhanced production of protochlorophyllide (Pchl)ide—a potent, natural  $^1\text{O}_2$  photosensitizer—(Meskauskiene *et al.*, 2001; Kaus *et al.*, 2012). When seedlings of the *flu* mutant are dark adapted, Pchl)ide accumulates in thylakoids. After a dark to light shift, a surge in  $^1\text{O}_2$  production occurs in chloroplasts, eventually leading to cell death. Direct photodamage by  $^1\text{O}_2$  takes place in chloroplasts of the *flu* mutant, but the ongoing events responsible for the cell death are genetically mediated by two plastid proteins EXECUTER1 (EX1) and EX2 (Lee *et al.*, 2007; Kim *et al.*, 2008; Kim and Apel, 2013). In other words, programmed cell death (PCD) is the default defence response that becomes active in the *flu* mutant. The role of the EX proteins has also been extended to wild-type plants, but the defence responses triggered seem different. Kim and co-workers (2012) proposed that EX-dependent signalling induces the formation of microlesions, but not the disintegration of chloroplasts, a finding that was interpreted as an acclimation response that enhances stress resistance in wild-type plants. Another *Arabidopsis* mutant where the  $^1\text{O}_2$ -mediated transcriptional responses have been investigated in detail is the double mutant *npq1lut2* that lacks zeaxanthin and lutein (Alboresi *et al.*, 2011). In *npq1lut2*, the accumulation of  $^1\text{O}_2$  is selectively enhanced; however, the transcriptional response does not initiate PCD, but rather the induction of genes whose function is to protect chloroplasts against the damaging effect of ROS. In two other mutants, where  $^1\text{O}_2$  production is selectively enhanced, *vte1 npq1*, a double mutant deficient in  $\alpha$ -tocopherol and zeaxanthin, and *chl1*, a mutant deficient in Chl *b*, high light (HL) irradiance led to a large increase of  $^1\text{O}_2$  that induced cell death as a consequence of a direct  $^1\text{O}_2$  cytotoxic effect (Triantaphylidès *et al.*, 2008), although it has recently been recognized that *chl1* can acclimate to  $^1\text{O}_2$  if exposed to mild light stress first (Ramel *et al.*, 2013).  $^1\text{O}_2$ -mediated transcriptional responses at HL irradiance were

also studied in wild-type plants and cell suspension cultures of *Arabidopsis* (González-Pérez *et al.*, 2011; Gutiérrez *et al.*, 2011; Ramel *et al.*, 2012b). HL stress revealed a set of transcripts that overlapped significantly with those listed in the *flu* mutant after the dark to light shift; however, in contrast to the *flu* mutant, an acclimatory response responsible for an increased tolerance against a more severe photooxidative stress was activated instead of PCD. It is worth noting that in wild-type plants  $^1\text{O}_2$  is produced at the heart of PSII (i.e. the PSII RC) and that the above studies provided evidence for  $^1\text{O}_2$  production in the PSII RC, for example non-enzymatic oxidation of the  $\beta$ -carotene of the PSII RC (Arellano *et al.*, 2011; Ramel *et al.*, 2012a) and photodamage to the D1 protein of the PSII RC (González-Pérez *et al.*, 2011). Furthermore, Ramel and co-workers (2012b) established that the differential transcriptional expression induced by  $\beta$ -carotene oxidation products was not mediated by the EX1 protein, and González-Pérez and co-workers (2011) showed no evidence for the up-regulation of *EDS1* encoding the ENHANCED DISEASE SUSCEPTIBILITY PROTEIN 1, also a gene key for understanding the  $^1\text{O}_2$ -mediated cell death response (Ochsenbein *et al.*, 2006). At present, it is not well established whether differences in the type of  $^1\text{O}_2$ -mediated transcriptional defence responses depend on the source of  $^1\text{O}_2$  within the chloroplasts (i.e. PSII RC, thylakoid-bound Chl precursors, infiltrated artificial photosensitizers, etc.), the intensity of the stress (time, concentration, etc.), or a combination of both, and questions about this issue have been raised in the past few years (Galvez-Valdivieso and Mullineaux, 2010; Alboresi *et al.*, 2011; Gutiérrez *et al.*, 2011; Ramel *et al.*, 2012a; Kim and Apel, 2013).

In an attempt to shed more light on the  $^1\text{O}_2$ -mediated cell death response in plants, Rose Bengal (RB)—a potent, artificial  $^1\text{O}_2$  photosensitizer—and dark-grown and light-grown *Arabidopsis* cells, the former containing proplastids, while the latter have functional chloroplasts, were used here. RB is an  $^1\text{O}_2$  elicitor that accumulates inside chloroplasts and has been used in several studies with the model alga *Chlamydomonas reinhardtii* to investigate the role of  $^1\text{O}_2$  in chloroplast to nucleus retrograde signalling and acclimatory responses to  $^1\text{O}_2$  stress (Leisinger *et al.*, 2001; Fischer *et al.*, 2007; Ledford *et al.*, 2007). In these studies, when cells were exposed to light, RB induced (in a similar fashion to HL stress) the transcription of *GLUTATHIONE PEROXIDASE HOMOLOGOUS* (*GPXH*), a gene with a key role in defending against photooxidative stress. In addition, constitutive overexpression of *GPXH* along with *GLUTATHIONE-S-TRANSFERASE 1* (*GSTSI*) was sufficient to augment  $^1\text{O}_2$  resistance. As with *C. reinhardtii*, *Arabidopsis* cell suspension cultures (ACSC) are an excellent plant model system to investigate  $^1\text{O}_2$  elicitor-mediated transcriptional defence responses and cell viability (or PCD) following chemical treatments. Quantification of PCD rates in cells *in vivo* can be rather difficult when infiltration of chemicals is required as the defence responses often occur in a small group of inaccessible cells buried in a bulk of surrounding healthy cells. Consequently, plant cell cultures can offer a more suitable means of investigating PCD due to their accessibility, reduced complexity, and uniformity

(McCabe and Leaver, 2000). Additionally, ACSC can also be grown in the dark without difficulty, which in turn represents another advantage when chloroplast-dependent physiological responses are under study (Doyle *et al.*, 2010).

In this study, ACSC were subjected to mild photooxidative conditions by adding RB at submicromolar concentrations to avoid side effects that were not directly related to  $^1\text{O}_2$  production. After the treatment with RB in the light,  $^1\text{O}_2$ -mediated PCD was only observed in *Arabidopsis* cells containing functional chloroplasts, and their transcriptional defence responses resembled those induced by  $^1\text{O}_2$  in the *flu* mutant. The comparative analysis of the transcriptional profiles of ACSC, treated with RB and  $\text{H}_2\text{O}_2$  in the light, provided some clues about transcripts with a key role in PCD. In order to test if cellular location was an important factor in responses induced by  $^1\text{O}_2$ , two other  $^1\text{O}_2$  elicitors were used in this study, Methylene Violet (MV) and Indigo Carmine (IC), which accumulate differently in plant cells (Hideg *et al.*, 2011). However, neither was able to induce any statistically significant change in transcript expression. The poor  $^1\text{O}_2$  quantum yield of MV and IC in comparison with that of RB (Kovács *et al.*, 2013) and, consequently, the need for higher concentrations to match equal levels of  $^1\text{O}_2$  production means that it cannot be established whether lack of change in transcript expression was due to different cellular location or possible limitations in their suitability as photosensitizers in studies where the matter of interest is the effect of  $^1\text{O}_2$  production by them, but not their own chemical toxicity.

## Materials and methods

### Growth conditions for *Arabidopsis* cell suspension cultures

Light-grown *Arabidopsis thaliana* L. (Columbia ecotype) cell suspension cultures (ACSC) were maintained in 200 ml of liquid growth medium (Jouanneau and Péaud-Lenoël, 1967; Axelos *et al.*, 1992) as previously described (González-Pérez *et al.*, 2011). For comparative purposes, *Arabidopsis* cells were also grown in the dark when cell viability was under investigation (see below).

### Cellular location of $^1\text{O}_2$ photosensitizers

Before proceeding with the photooxidative treatments, the location of RB, MV, and IC inside the cells of *Arabidopsis* was investigated by confocal microscopy. The absorption and emission spectral features of these three  $^1\text{O}_2$  photosensitizers have been previously described (Morrison *et al.*, 1997; Roberts *et al.*, 1998; Chang *et al.*, 2008). A 1 ml aliquot of ACSC, previously treated with 1 mM of each photosensitizer for 10 min in the dark, was centrifuged at 120 *g* for 3 min at 4 °C, washed three times with 2 ml of 2.7 mM KCl, 147.3 mM NaCl, and 10 mM sodium phosphate [phosphate-buffered saline (PBS)] pH 7.4, and finally resuspended in 1 ml of 10 mM PBS pH 7.4. The fluorescence emission was then monitored with a confocal microscope (model DM IRB; Leica Microsystems) following excitation of the three photosensitizers and Chl at 543 nm and 633 nm with an argon laser and a Triplet Dichroic 488/543/633 excitation beam splitter. The fluorescence emission of the selected photosensitizers and chloroplast Chl-binding proteins was collected at 560–620 nm and at 680–700 nm, respectively.

### Photooxidative treatments

Nine-day-old cultures with a cell density of ~125–150 mg ml<sup>-1</sup> fresh weight were split into two batches, acclimatized before the chemical treatment for 2 h in the dark or continuous illumination at 150  $\mu\text{E m}^{-2} \text{s}^{-1}$  at 24 °C with gentle agitation, and then treated with the photosensitizers at 0.5  $\mu\text{M}$  and 5  $\mu\text{M}$ . Cultures were also treated with 0.5 mM and 5 mM  $\text{H}_2\text{O}_2$  for comparative purposes. After 30 min of chemical treatment, samples were collected, filtered, frozen in liquid nitrogen, and stored at -80 °C until further analysis. Light and dark treatments were carried out simultaneously, performed in duplicate, and replicated three times. Control cultures were also exposed to the same conditions either at 150  $\mu\text{E m}^{-2} \text{s}^{-1}$  or in the dark.

### Photosynthetic activity

In order to determine the photosynthetic activity, the rate of oxygen production was measured polarographically using a Chlorolab 2 system (Hansatech Instruments, Norfolk, UK) at 20 °C. Samples containing equal amounts of fresh weight of cells were incubated in the dark for 1 min before a 300  $\mu\text{E m}^{-2} \text{s}^{-1}$  light-emitting diode source was switched on. The contribution of mitochondrial respiration to the rate of oxygen production was subtracted when the same experiment was repeated with a fresh sample in the dark. The photosynthetic oxygen evolution rate was expressed as  $\mu\text{M min}^{-1} \text{g}^{-1}$  of fresh weight of cells and the percentage of photosynthetic activity was referred to control cells. All the rate measurements were performed in triplicate.

### Protein oxidation analysis

The oxidative damage to the D1 protein (i.e. PsbA) of the PSII RC and general protein oxidation in ACSC were evaluated by western blot. Proteins were extracted from ~500 mg of frozen cells as previously described (González-Pérez *et al.*, 2011). Protein samples were subjected to 6 M urea, 12% (w/v) SDS-PAGE and transferred overnight to nitrocellulose membranes. Nitrocellulose membranes were stained with Ponceau S solution (Applichem, Darmstadt, Germany) for 1 min to visualize the protein transfer and then destained with water. The nitrocellulose membranes were blocked with 1% (w/v) bovine serum albumin (BSA) in 0.6% (w/v) Tween-20, 150 mM NaCl, 50 mM TRIS-HCl, pH 7.4 (T-TBS). For the immunodetection of PsbA, the nitrocellulose membranes were incubated for 1 h at 37 °C with polyclonal antibodies against the C-terminal region of PsbA (Agrisera, Vännäs, Sweden) using a dilution of 1:5000 in T-TBS. After extensive washing in T-TBS, the immunocomplexed membranes were probed for 1 h at 37 °C with an anti-rabbit, peroxidase-linked secondary antibody using a dilution of 1:2000 in T-TBS, and were washed with T-TBS once again. The immunodetection of PsbA was visualized with the 3,3'-diaminobenzidine tetrahydrochloride (DAB) substrate kit (Pierce Biotechnology, Rockford, IL, USA) or with luminol (Bio-Rad Laboratories, Hercules, CA, USA). To detect general protein oxidation, the extracted proteins were chemically modified with 2,4-dinitrophenylhydrazine (DNPH) and the corresponding carbonylated proteins were identified following the steps described in the OxyBlot protein oxidation detection kit from Millipore (Millipore, Billerica, MA, USA), and visualized with luminol (Bio-Rad Laboratories).

### Indirect detection of $^1\text{O}_2$

The production of  $^1\text{O}_2$  by RB was analysed in ACSC that were first broken in water or deuterium oxide ( $\text{D}_2\text{O}$ ), and then treated with RB at a concentration of 5  $\mu\text{M}$  under continuous illumination at 150  $\mu\text{E m}^{-2} \text{s}^{-1}$  or in the dark for 30 min. After the treatment, the electrophoretic migration of the band of the D1/D2 heterodimer was inspected by western blot analysis, as described above, to evaluate  $^1\text{O}_2$  production qualitatively (Lupinkova and Komenda, 2004; Arellano *et al.*, 2011).



### Measuring cell death levels

After the (photo)oxidative treatments, dark-grown or light-grown cells were diluted 20 times into fresh culture medium and left in the dark for 24 h to allow the progression of cell death-associated morphological changes. The vital stain fluorescein diacetate (FDA) was applied to ACSC to assess their cell viability. Cells were subsequently scored as alive, dead via necrosis, or dead via PCD using a UV-equipped light microscope according to the methods previously described (McCabe and Leaver, 2000; Reape *et al.*, 2008; Doyle *et al.*, 2010). Under UV radiation, a bright green fluorescence is observed in viable cells with an intact plasma membrane. Dead cells that displayed the hallmark PCD morphology consisting of a condensed protoplast, which had retracted away from the cell wall, were scored as dead via PCD. FDA-stained cells that did not exhibit bright green fluorescence or a retracted cell protoplast were scored as necrotic.

### Target transcripts to evaluate early ROS-mediated responses

Target transcripts to monitor the responses of ACSC under the assayed oxidative stress conditions were selected from lists containing ROS markers specifically up-regulated by  $^1\text{O}_2$  and  $\text{H}_2\text{O}_2$  (op den Camp *et al.*, 2003; Gadjev *et al.*, 2006; Laloi *et al.*, 2007). The *DEFENSIN-LIKE FAMILY (DEF)* transcript named At2g43510 was chosen as a marker for general oxidative stress (Gadjev *et al.*, 2006), whereas the *PROFILIN FAMILY (PROF)* transcript named At2g19760 was selected as a housekeeping marker for internal reference (Laloi *et al.*, 2007). The primers designed to amplify the selected transcripts are given in Supplementary Table S1 available at JXB online.

### RNA isolation and RT-PCR analysis

RNA isolation and relative quantification of mRNA expression were performed as previously described (Livak and Schmittgen, 2001; Chomczynski and Sacchi, 2006; González-Pérez *et al.*, 2011). Other intermediate steps such as contaminating genomic DNA elimination, RNA reverse transcription, and reverse transcription-PCR (RT-PCR) runs are as previously described (González-Pérez *et al.*, 2011). Expression levels were normalized using the housekeeping gene *PROF* (At2g19760).

### Microarray experiments and data analysis

Transcriptional analyses were performed using Affymetrix GeneChip Arabidopsis genome ATH1 arrays (Affymetrix Inc., Santa Clara, CA, USA). The method to verify the quality of total, DNA-free RNA, and other technical details describing the microarray experiments are as previously described (González-Pérez *et al.*, 2011). Experiments were performed using three biological replicates. Principal components analysis (PCA) was used to explore the variability between the replicates (Johnson and Wichern, 1998). Data from microarrays were standardized using quantile normalization and the robust multi-array average (RMA) method (Bolstad *et al.*, 2003). Differential transcript expression was carried out using the Limma package (Smyth, 2004) available from Bioconductor (<http://www.bioconductor.org/>). Multiple testing adjustments of *P*-values were carried out according to the Benjamini and Hochberg methodology (Benjamini and Hochberg, 1995). A volcano plot was used to visualize transcripts with statistically significant differential expression (Cui and Churchill, 2003). Significantly over- or under-represented gene ontology (GO)/biological process terms were obtained using FatiScan from the Babelomics suite (<http://babelomics.bioinfo.cipf.es/>) as described previously (Al-Shahrour *et al.*, 2004, 2007; Medina *et al.*, 2010). Multiple testing adjustment of *P*-values was then carried out according to the false discovery rate (FDR) method (Benjamini and Hochberg, 1995; Benjamini and Yekutieli, 2001). GO terms were annotated from the Ensembl's 56 (TAIR 10) release (<http://www.ensembl.org/>).

### Validation of microarray experiments

In order to validate the microarray experiments, several transcripts with statistically significant differential expression were selected from the list included in Supplementary Tables S2 and S3 at JXB online. The selected transcripts and their corresponding primers are given in Supplementary Table S4.

### Co-regulation analysis

Data from the microarray experiments were clustered together with expression data from previously selected key treatments in *Arabidopsis* based on comparative analysis using Genevestigator (Hruz *et al.*, 2008). Only transcripts presenting signal  $\text{Log}_2$  ratios  $\geq 1$  (induction) or  $\leq -1$  (repression) were considered for analysis. The hierarchical clustering analysis was then carried out using R statistical software, version 2.12.1 (<http://www.r-project.org>). Pearson's correlation analysis was performed to evaluate the linear relationship between experimental treatments (Quinn and Keough, 2002). The raw microarray data have been deposited in the GEO database under the accession number GSE43551.

## Results

### Cellular location of the $^1\text{O}_2$ photosensitizers in ACSC


The cellular location of the  $^1\text{O}_2$  photosensitizers RB, MV, and IC was investigated on the basis of their fluorescence using a confocal microscope (Fig. 1). Additionally, red Chl autofluorescence was used to identify the chloroplasts inside the cells (Fig. 1I–L). The incubation with each photosensitizer was performed in the dark for only 10 min to avoid unnecessary cellular damage. The *Arabidopsis* cells treated with 1 mM IC did not exhibit fluorescence emission (Fig. 1F), suggesting that this dye barely permeates the cell or it was not retained within the cell after washing with 10 mM PBS pH 7.4. In contrast to IC, the *Arabidopsis* cells treated with 1 mM MV or 1 mM RB showed a distinct green fluorescence around and inside the cells that clearly indicated that these photosensitizers were both able to cross the plasma membrane and to be partly retained on the cell boundary (i.e. cell wall and plasma membrane), and other cellular compartments (Fig. 1G, H). In spite of the ability of MV and RB to cross the cell boundary, a closer inspection of the fluorescence images revealed a significant difference between the dyes. MV accumulated in the chloroplast envelope and showed a ring-like fluorescence emission around these organelles, while the red Chl autofluorescence was still discernible (Fig. 1G, O), indicating that MV apparently did not reach the chloroplast interior. In contrast, the distinctive red Chl autofluorescence was blurred by the green fluorescence emission of RB (Fig. 1H, P), showing clearly that this dye could indeed cross the chloroplast envelope and accumulate inside this organelle.

### Protein oxidation in ACSC as marker for photooxidative stress level


Experimental conditions were chosen to induce mild photooxidative stress in *Arabidopsis* cells in order to investigate defence responses that would be activated in a series of scenarios where the partitioning of  $^1\text{O}_2$  between the surrounding

medium, the cells, and cellular organelles was imposed by the chemical nature of the  $^1\text{O}_2$  photosensitizers. Additionally, *Arabidopsis* cells were exposed to  $\text{H}_2\text{O}_2$  for comparative purposes, and its concentration was adjusted to induce a level of oxidative stress similar to that caused by the three  $^1\text{O}_2$  elicitors. In order to avoid the superimposition of other signaling pathways that could become activated by severe damage,

the lowest possible concentrations of RB, MV, IC, and  $\text{H}_2\text{O}_2$  were chosen where oxidative damage was perceived in the first 30 min of the light treatment. The level of carbonylated proteins was measured as a marker for oxidative stress in the cellular proteome of ACSC. Figure 2 shows the changes in the oxidation state of the cell proteome after the 30 min treatment with two different concentrations for RB,



**Fig. 1.** Representative confocal micrographs illustrating the location of IC, MV, and RB at a concentration of 1 mM in ACSC. From top to bottom the following panels are represented: bright field images (A–D),  $^1\text{O}_2$  elicitor fluorescence images at 560–620 nm (E–H), red Chl autofluorescence images at 680–700 nm (I–L), merged Chl and  $^1\text{O}_2$  elicitor fluorescence images (M–P), and merged bright field and fluorescence images (Q–T). The  $^1\text{O}_2$  elicitor and red Chl fluorescence emissions were collected after excitation at 543 nm and 633 nm with an argon laser. Arrows in G, H, O, and P point to key features (see text for further details). Scale bars=50  $\mu\text{m}$ .



**Fig. 2.** Protein oxidation analysis OxyBlot of ACSC under (photo)oxidative stress conditions. Control (A), RB (B), MV (C), IC (D), and  $\text{H}_2\text{O}_2$  (E). ACSC were treated for 30 min under continuous illumination at  $150 \mu\text{E m}^{-2} \text{s}^{-1}$  (L) or in the dark (D) at two different concentrations for each photosensitizer (0.5  $\mu\text{M}$  and 5  $\mu\text{M}$ ) and  $\text{H}_2\text{O}_2$  (0.5 mM and 5 mM). Each lane was loaded with 30  $\mu\text{g}$  of protein.

MV, IC (i.e. 0.5  $\mu\text{M}$  and 5  $\mu\text{M}$ ), and for  $\text{H}_2\text{O}_2$  (i.e. 0.5 mM and 5 mM). A light treatment of ACSC with the photosensitizers at a concentration of 10  $\mu\text{M}$  after 30 min showed clear evidence of cellular lysis and a loss of cellular material after filtration, particularly in the case of RB. Additionally, homogenous light penetration in the culture medium was hampered at concentrations  $>10 \mu\text{M}$  due to the high molar absorptivity of the photosensitizers. Consequently, the concentration of 5  $\mu\text{M}$  for each photosensitizer was considered as the upper limit for this photooxidative analysis. In the absence of the photosensitizers or  $\text{H}_2\text{O}_2$ , the OxyBlot experiments showed the immunodetection of a basal level of protein oxidation that did not differ appreciably between light and dark conditions after 30 min (Fig. 2). This was not the case when the photosensitizers were added to the medium. The light treatments of ACSC in the presence of RB, MV, and IC at 5  $\mu\text{M}$  revealed an evident increase in protein oxidation in their respective lanes that was not accompanied by similar changes in the dark treatments for the same concentration, showing a clear light-dependent effect on the protein oxidation of ACSC. The changes in protein oxidation of ACSC at 0.5  $\mu\text{M}$  were subtle, but still discernible, for each photosensitizer when the respective treatments in light and dark conditions were compared. Therefore, 0.5  $\mu\text{M}$  for each photosensitizer was considered high enough to induce mild photooxidative stress. When compared with control ACSC, the treatment of ACSC with  $\text{H}_2\text{O}_2$  enhanced protein oxidation, but it was not light dependent. On the contrary, protein oxidation became more apparent in the dark, although this finding was not investigated further.

#### RT-PCR analysis of selected transcripts responding to photooxidative stress

Changes in the expression profile for some specific markers for  $^1\text{O}_2$  and  $\text{H}_2\text{O}_2$  were investigated after a 30 min treatment with RB, MV, and IC at 0.5  $\mu\text{M}$ , and  $\text{H}_2\text{O}_2$  at 500  $\mu\text{M}$ . For the sake of clarity, the full name of the chosen specific markers and their respective locus identifiers are given in Supplementary Table S1 at *JXB* online. As shown in Supplementary Table S5, significant changes were observed in the expression profile of some selected markers after the treatments with RB. However, no significant changes in the expression profile of any of the selected ROS markers were observed after the treatments with IC and MV, a result that was further confirmed in the whole-genome transcriptional profiling analysis (see below). When ACSC were exposed to RB, the specific  $^1\text{O}_2$  markers *NOD* and *SZF1* were significantly up-regulated, whereas the specific  $\text{H}_2\text{O}_2$  marker *SUBT* was found to be down-regulated. The same ROS markers (i.e. *NOD*, *SZF1*, *SUBT*, and *F9D12*) did not show any significant change under dark conditions during the 30 min treatment with 0.5  $\mu\text{M}$  RB. No significant changes in the expression profile of the general oxidative stress marker (*DEF*) were observed in the presence of any of the  $^1\text{O}_2$  elicitors. In contrast to this, the 30 min treatment with  $\text{H}_2\text{O}_2$  resulted in slight up-regulation of *DEF*, but not of any of the selected  $\text{H}_2\text{O}_2$  markers, confirming that 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$

also induced mild oxidative stress in ACSC under the experimental conditions used here.

#### Limma analyses of $^1\text{O}_2$ - and $\text{H}_2\text{O}_2$ -mediated transcriptional responses of ACSC

The  $^1\text{O}_2$ - and  $\text{H}_2\text{O}_2$ -mediated transcriptional responses of ACSC were further characterized using Affymetrix GeneChip *Arabidopsis* genome ATH1 arrays. The microarray experiments were designed and classified as follows: control (C1, C2, and C3), RB at 0.5  $\mu\text{M}$  (RB1, RB2, and RB3), MV at 0.5  $\mu\text{M}$  (MV1, MV2, and MV3), IC at 0.5  $\mu\text{M}$  (IC1, IC2, and IC3), and  $\text{H}_2\text{O}_2$  at 500  $\mu\text{M}$  (HP1, HP2, and HP3); where the numbers 1–3 represent the number of biological replicates. After data normalization, the Limma package identified a total of 1705 transcripts which were differentially expressed in ACSC exposed to RB with an adjusted *P*-value  $<0.05$  (Supplementary Table S2 at *JXB* online). Out of a total 1705 transcripts, 314 had a fold change  $>1$  ( $\text{Log}_2$ ), whereas 171 exhibited a fold change  $<-1$ . Surprisingly, no transcripts with statistically significant differential expression (adjusted *P*-value  $<0.05$ ) were found in cell cultures treated with either MV or IC. In comparison with the RB treatment, the treatment of ACSC with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  induced a mild transcriptional response characterized by the observation that both the number of transcripts differentially expressed (i.e. 568, adjusted *P*-value  $<0.05$ ) and the number of transcripts with a fold change  $<-1$  or  $>1$  (i.e. 270) were lower than in the RB treatment (Supplementary Table S3). Volcano plots are shown in Supplementary Fig. S1 to better summarize the experimental conditions that yielded statistically significant changes in transcript expression. It is worth noting that 32 out of the 314 up-regulated transcripts (and four out of the 171 down-regulated transcripts) with a statistically significant fold change  $|\text{Log}_2|>1$  after treatment with 0.5  $\mu\text{M}$  RB were included in the lists of 296 up-regulated and 29 down-regulated transcripts described as specific markers for  $^1\text{O}_2$  according to the transcriptional profile analyses performed by Gadjev *et al.*, (2006). Approximately 45% and 30% of the transcripts with up- and down-regulation after the treatment with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  were also present in the lists of up- and down-regulated transcripts after treatment with 0.5  $\mu\text{M}$  RB. Surprisingly, the treatment with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  induced the expression ( $|\text{Log}_2|>1$ ) of only three transcripts defined as specific markers for  $\text{H}_2\text{O}_2$ , but 19 transcripts defined as specific markers for  $^1\text{O}_2$  (Gadjev *et al.*, 2006) (Supplementary Table S6).

RT-PCR and Limma analyses were in line with respect to the treatments with 0.5  $\mu\text{M}$  MV and 0.5  $\mu\text{M}$  IC, and both confirmed that there were no statistically significant changes in the transcriptional profile of ACSC after these treatments. It was first hypothesized that the different cellular location of the three photosensitizers could explain the divergent changes in the transcriptional profile. However, it was also found reasonable that the differences were based on the inherent quantum yields of  $^1\text{O}_2$  for RB, MV, and IC, which are in a ratio of 0.75:0.37:0.095 (Kovacs *et al.*, 2013). Although the most obvious thing to do was to adjust the MV and IC



concentrations to reach  $^1\text{O}_2$  production levels similar to RB, several factors such as cellular lysis, non-homogeneous light penetration, and a direct cytotoxic effect by the photosensitizers discouraged the use of higher concentrations. Therefore, it could not be unambiguously determined whether cellular location was the sole reason why no significant changes in the transcriptional profile of ACSC were observed after the treatments with IC and MV, and, consequently, the study was centred on the treatments with RB and  $\text{H}_2\text{O}_2$ .

#### Microarray data validation by RT-PCR

The microarray data of the 30 min treatments with  $0.5 \mu\text{M}$  RB or  $500 \mu\text{M}$   $\text{H}_2\text{O}_2$  were confirmed by RT-PCR. All the transcripts selected for validation (Supplementary Table S4 at *JXB* online) showed similar fold changes in expression when they were analysed by both techniques (Supplementary Fig. S2).

#### GO process terms associated with response to stimuli after (photo)oxidative treatments


In an attempt to identify GO/biological process terms that became over-represented after the treatments with RB and  $\text{H}_2\text{O}_2$ , functional enrichment was carried out using FatiScan from the Babelomics suite (<http://babelomics.bioinfo.cipf.es>). Briefly, this tool makes use of the complete list of genes available in the ATH1 arrays (i.e.  $\sim 24\,000$ ) and detects significantly up- or down-regulated blocks of functionally related transcripts ordered by differential expression after the treatment, no matter whether the differential transcriptional expression is found to be statistically significant or not. The treatment with  $0.5 \mu\text{M}$  RB exhibited an extensive list of over-represented biological processes (42 GO terms) that were associated with

responses to (i) stimuli (i.e. 25 out of 42; defence and immune responses, responses to abiotic, biotic, organic substance, etc.) and (ii) metabolic, cellular, and developmental processes (i.e. 17 out of 42; nitrogen, nucleic acid, and protein metabolic processes, transcription, protein folding and modification, cell cycle, PCD, apoptosis, etc.) (Supplementary Table S7 at *JXB* online). In the presence of  $500 \mu\text{M}$   $\text{H}_2\text{O}_2$ , 46 GO terms were over-represented with statistical significance and they could be divided into GO terms ascribed to responses to (i) stimuli (i.e. 20 out of 46) and (ii) metabolic, cellular, and developmental processes (i.e. 27 out of 46) (Supplementary Table S8). Sixteen out of the 20 over-represented defence processes for the treatment with  $\text{H}_2\text{O}_2$  were common with the RB treatment (Supplementary Tables S7 and S8). Other GO/defence process terms for the  $\text{H}_2\text{O}_2$  treatment were associated with metal ion responses.

#### RB and $\text{H}_2\text{O}_2$ inhibit the photosynthetic activity of ACSC

The photosynthetic oxygen evolution rate of ACSC was examined in the presence of  $0.5 \mu\text{M}$  RB and  $500 \mu\text{M}$   $\text{H}_2\text{O}_2$  (Fig. 3A). It decreased to  $\sim 65\%$  of the initial value under control conditions when RB was present, suggesting that the  $^1\text{O}_2$  produced by RB inside chloroplasts could inhibit the photosynthetic activity of ACSC. The treatment with  $500 \mu\text{M}$   $\text{H}_2\text{O}_2$  also brought a slight inhibition of the photosynthetic activity of ACSC, indicating that exogenous  $\text{H}_2\text{O}_2$  easily permeated the plasma membrane and entered chloroplasts.

In order to establish whether the presence of  $^1\text{O}_2$  and  $\text{H}_2\text{O}_2$  could damage the D1 protein of PSII in ACSC, cultured cells were subjected to a western blot analysis using an antibody raised against the D1 protein (Fig. 3B). In control conditions,



**Fig. 3.** (A) Oxygen evolution rates of ACSC under control conditions (C) and after the 30 min treatments with  $0.5 \mu\text{M}$  RB and  $500 \mu\text{M}$   $\text{H}_2\text{O}_2$ , \**P*-value  $< 0.05$ . (B and C) Western blot analysis of the D1 protein of PSII after the 30 min treatment with  $0.5 \mu\text{M}$  RB and  $500 \mu\text{M}$   $\text{H}_2\text{O}_2$  (HP) under continuous illumination (B) and in the dark (C). Lane 1, control (C); lane 2, RB; lane 3, HP. Each lane was loaded with  $70 \mu\text{g}$  of protein. The upper, middle, and lower band show the D1/D2 heterodimer of PSII, the adduct between the monomeric D1 protein and the  $\alpha$ -subunit of the cytochrome *b559* (D1/PsbE), and the monomeric D1 (D1), respectively. (D) Western blot analysis of the D2/D1 heterodimer of PSII after the 30 min treatment of broken ACSC with  $5 \mu\text{M}$  RB under continuous illumination at  $150 \mu\text{E m}^{-2} \text{s}^{-1}$  (L) or in the dark (D) in water or deuterium oxide ( $\text{D}_2\text{O}$ ). Each lane was loaded with  $30 \mu\text{g}$  of protein.


three bands were distinguished; the lower and more intense band corresponded to the monomeric D1 protein of the PSII RC, whereas the upper band corresponded to the D1/D2 heterodimer of the PSII RC. The intermediate band, which is often observed to react against the D1 antibody, was a faint band that corresponded to an adduct between the D1 protein and the  $\alpha$ -subunit of cytochrome  $b_{559}$  of PSII (Arellano *et al.*, 2011, and references therein). The intensity of the band ascribed to the monomeric D1 protein was lower in the presence of 0.5  $\mu\text{M}$  RB, but it does not change in the treatment with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  (Fig. 3B). As a control experiment, the incubation with RB and  $\text{H}_2\text{O}_2$  was also performed in the dark, and the results showed no damage to the D1 protein (Fig. 3C). Neither significant damage to the D1 protein nor a substantial decrease in the photosynthetic oxygen evolution rate were to be expected in the presence of 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  on the basis of the study by Miyao *et al.* (1995), who reported a more visible and progressive damage to the D1 protein in the range of millimolar concentrations of  $\text{H}_2\text{O}_2$ . However, this does not exclude that other photosynthetic complexes (i.e. PSI or Rubisco) could be more sensitive to  $\text{H}_2\text{O}_2$  inhibition (Nakano *et al.*, 2006).

To demonstrate qualitatively that RB produced  $^1\text{O}_2$  under the present experimental conditions, broken ACSC were treated with this photosensitizer at a concentration of 5  $\mu\text{M}$  in both water and  $\text{D}_2\text{O}$ , where the lifetime of singlet oxygen notably varies from  $\sim 4$   $\mu\text{s}$  to  $\sim 70$   $\mu\text{s}$ . It is known that the D1/D2 heterodimer exhibits a shift in its electrophoretic migration

due to changes in protein conformation after oxidation of amino acid residues prone to react with  $^1\text{O}_2$  (Lupinkova and Komenda, 2004; Arellano *et al.*, 2011). Figure 3D showed that 5  $\mu\text{M}$  RB induced a shift of the immunodetected band of the D1/D2 heterodimer toward the cathodic end of the gel together with a slight broadening after the light treatment. The shift was more prominent when the broken ACSC were exposed to the 30 min light treatment in  $\text{D}_2\text{O}$ , where the lifetime for  $^1\text{O}_2$  was longer and, consequently, the changes in the conformation of the D1/D2 heterodimer induced by  $^1\text{O}_2$  were more acute.


#### *RB, but not $\text{H}_2\text{O}_2$ , affects PCD rates in light-grown ACSC*

The production of ROS (and particularly  $^1\text{O}_2$ ) in chloroplasts has been demonstrated to play a key role in PCD regulation (Doyle *et al.*, 2010; Kim *et al.*, 2012). Therefore, the two treatments with RB or  $\text{H}_2\text{O}_2$ —both compounds able to enter chloroplasts—could affect the viability of ACSC and induce PCD. In order to determine if this was the case, visible morphological PCD changes, together with FDA fluorescence, were investigated. Figure 4 shows four representative bright field images of light-grown cells under control conditions, after treatment with 0.5  $\mu\text{M}$  RB or 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  for 30 min under continuous illumination at 150  $\mu\text{E m}^{-2} \text{s}^{-1}$ , and after exposure to HL (i.e. 1800  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) for 90 min. Under control conditions, light-grown cells exhibited the standard morphology (i.e. excessive hydration in cellular



**Fig. 4.** Representative field images showing changes in the morphology of light-grown *Arabidopsis* cells under control conditions (A), after the 30 min treatment with 0.5  $\mu\text{M}$  RB (B) and 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  (C), and after the 90 min treatment at 1800  $\mu\text{E m}^{-2} \text{s}^{-1}$  (D). Bright green fluorescence of fluorescein diacetate following excitation with UV radiation in A, C, and D indicates that *Arabidopsis* cells remain viable after the stress treatment. Arrows in B and D illustrate some of the gaps between the cell wall and the plasma membrane after the induction of PCD. Scale bars=50  $\mu\text{m}$ .





**Fig. 5.** PCD in light-grown and dark-grown *Arabidopsis* cells subjected to 30 min chemical treatments at  $150 \mu\text{E m}^{-2} \text{s}^{-1}$  or in the dark. Black bar,  $0.5 \mu\text{M RB}$ ; grey bar,  $500 \mu\text{M H}_2\text{O}_2$ ; white bar, control samples. \* $P$ -value  $< 0.05$ .

cultures) and were largely viable (12–15% PCD; Fig. 5), as demonstrated by the FDA fluorescence. In contrast to this, a clear gap between the cell wall and the plasma membrane was discernible in many of the light-grown cells treated with  $0.5 \mu\text{M RB}$ , and bright green fluorescence could barely be detected in the cells stained with FDA (Fig. 4), indicating that PCD was greatly induced in light-grown ACSC after the RB treatment (79%; Fig. 5). The condensation of the cell protoplast could not be observed when light-grown cells were subjected to  $500 \mu\text{M H}_2\text{O}_2$ , and bright green fluorescence was detected in the presence of FDA (Fig. 4). PCD was 14% (Fig. 5), indicating that the assayed concentration of  $\text{H}_2\text{O}_2$  did not induce PCD significantly under these experimental conditions and that higher  $\text{H}_2\text{O}_2$  concentrations are required for the activation of PCD (Desikan *et al.*, 1998, 2001). Likewise, cell viability of light-grown cells exposed to HL was examined to explore whether  $^1\text{O}_2$  production by PSII was capable of activating PCD. HL treatment at  $1800 \mu\text{E m}^{-2} \text{s}^{-1}$  for 30 min did not affect cell viability (González-Pérez *et al.*, 2011) and only HL treatments for (or beyond) 90 min resulted in morphological changes associated with PCD (Fig. 4). No significant effects were observed on PCD rates when light-grown ACSC were subjected to chemical treatments with  $0.5 \mu\text{M RB}$  or  $500 \mu\text{M H}_2\text{O}_2$  for 30 min in the dark (Fig. 5).

Further experiments were performed with dark-grown cells containing proplastids instead of functional chloroplasts (Doyle *et al.*, 2010), and cells were subjected to chemical treatments with  $0.5 \mu\text{M RB}$  or  $500 \mu\text{M H}_2\text{O}_2$  for 30 min under continuous illumination at  $150 \mu\text{E m}^{-2} \text{s}^{-1}$  or in the dark. The results showed that the viability of the chemically treated cells was similar to that of control cells for dark-grown cells; there was no effect whether the chemical treatment was with  $0.5 \mu\text{M RB}$  or  $500 \mu\text{M H}_2\text{O}_2$ , or whether the cells were exposed to light or remained in the dark during the RB or  $\text{H}_2\text{O}_2$  treatments (Fig. 5). Considered together, these results showed that the presence of developed chloroplasts

and illumination were essential for the activation of PCD in ACSC treated with  $0.5 \mu\text{M RB}$ .

#### Co-regulation of $^1\text{O}_2$ -producing mutants and HL treatments with RB-treated ACSC

Over the last 14 years, analysis of the transcriptional profile of the *flu* mutant family has shed much light on  $^1\text{O}_2$ -mediated defence responses in plants and has also provided lists of specific transcriptional markers and *cis*-regulatory elements (Meskauskiene *et al.*, 2001; op den Camp *et al.*, 2003; Gadjev *et al.*, 2006; Petrov *et al.*, 2012). The *flu* mutant family is characterized by the high production of  $^1\text{O}_2$  after the dark–light shift in response to the accumulation of Pchl<sub>ide</sub> in thylakoids. In contrast to the *flu* mutant, high levels of  $^1\text{O}_2$  are produced in response to excess excitation energy in PSII of wild-type plants when photosynthetic activity is inhibited (Mullineaux and Baker, 2010). It is well established that  $^1\text{O}_2$  is the major ROS generated under HL conditions (Triantaphylidès *et al.*, 2008; Triantaphylidès and Havaux, 2009) and, consequently, the activation of defence responses under HL conditions ought to share a set of  $^1\text{O}_2$ -responsive transcripts with the *flu* mutant family. GENEVESTIGATOR can be used to perform comparative genome-wide analyses under different experimental conditions (Hruz *et al.*, 2008), and here it has been used to identify light treatments or *Arabidopsis* mutants that exhibit a transcriptional expression pattern similar to that of ACSC treated with RB. Six experimental conditions are described in GENEVESTIGATOR as HL studies, although the light irradiance used in those studies ranged from  $400 \mu\text{E m}^{-2} \text{s}^{-1}$  to  $1800 \mu\text{E m}^{-2} \text{s}^{-1}$  (Vandenabeele *et al.*, 2004; Walters *et al.*, 2004; Vanderauwera *et al.*, 2005; Kleine *et al.*, 2007; Rossel *et al.*, 2007; Caldana *et al.*, 2011; González-Pérez *et al.*, 2011). Additionally, the GENEVESTIGATOR database contains the transcriptional profiling of *Arabidopsis* after perturbation with UV radiation, a ‘light’ perturbation also known to lead to the production of  $^1\text{O}_2$  and other ROS at high doses (Brosche *et al.*, 2002; Ulm *et al.*, 2004; Hideg *et al.*, 2013). Among the *Arabidopsis* mutants described to produce  $^1\text{O}_2$ , the database of GENEVESTIGATOR only includes the *flu* mutant family (Laloi *et al.*, 2007), but not others. The CEL files of the double mutant *npq1lut2* were kindly provided by Dr Alboresi and co-workers (Alboresi *et al.*, 2011) and were included in the genome-wide analysis. The study by Ramel and co-workers (2012b, 2013) could not be included in the co-regulation analysis because these authors used CATMAv5 instead of Affymetrix microarrays (Ramel *et al.*, 2012a, b), and only a manual inspection was performed with their microarray data (see the Discussion). Using a list of 400 transcripts of the  $0.5 \mu\text{M RB}$  treatment with  $|\text{Log}_2| > 1$ , it was found that the biclustering tool of GENEVESTIGATOR identified a group of transcripts exhibiting an expression pattern similar to that described in the *flu* mutant family. Additionally, other treatments such as HL irradiance beyond  $1400 \mu\text{E m}^{-2} \text{s}^{-1}$  or UV radiation also proved to have a similar expression pattern. In contrast, HL treatments below  $1400 \mu\text{E m}^{-2} \text{s}^{-1}$  showed a poor correlation. In a more robust analysis, the total number of 485 transcripts with significant differential

expression (with  $|\text{Log}_2| > 1$ ) in the treatment with 0.5  $\mu\text{M}$  RB was subjected to hierarchical clustering in order to identify both experimental conditions with similar differential transcript expression and clusters of transcripts that could be rich in specific markers for  $^1\text{O}_2$ .


In the first instance, it is worth noting that the *hy5* (*long hypocotyl5* transcription factor mutant of *Arabidopsis*) and *npq1lut2* mutants were subjected to light irradiance of 1000  $\mu\text{E m}^{-2} \text{s}^{-1}$  (Kleine *et al.*, 2007; Alboresi *et al.*, 2011) and exhibited a poor correlation with the other selected experimental systems (Supplementary Table S9 at *JXB* online); they clustered together with catalase-deficient (CAT2HP1) plants (Fig. 6). CAT2HP1 plants in HL accumulate  $\text{H}_2\text{O}_2$  (Vanderauwera *et al.*, 2005) and exhibit a distinctive transcriptional profile when compared with other experimental systems where  $^1\text{O}_2$ , ozone ( $\text{O}_3$ ), or superoxide radical ( $\text{O}_2^-$ ) accumulates (Gadjev *et al.*, 2006). The other set of experiments clustered and, in particular, the transcriptional profile of the treatment with 0.5  $\mu\text{M}$  RB showed the highest correlation with the *flu* family mutants (Supplementary Table S9).

In the second instance, two main clusters of transcripts were observed and named A and B (Fig. 6). Cluster A consisted of 242 transcripts and was characterized by the presence of the largest number of specific markers for  $^1\text{O}_2$  (i.e. 27 out of 36). The value of 3.42 for the odds ratio [i.e.  $27 \times (242 - 9) / 9 \times (243 - 27)$ ] indicated that the enrichment of specific markers for  $^1\text{O}_2$  in cluster A was statistically significant ( $P$ -value of  $1.5 \times 10^{-3}$ ). Cluster B with 243 transcripts included all the down-regulated transcripts with a fold change  $< 1$  in the treatment with 0.5  $\mu\text{M}$  RB and it did not show any significant enrichment in specific markers for any type of ROS. Table 1 summarizes the list of specific markers for ROS identified in cluster A. It is worth noting that six out of the 27 specific markers for  $^1\text{O}_2$  identified in cluster A were also found in the list of the 229 up-regulated transcripts with a fold change  $> 1$  in the treatment with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  (Supplementary Table S6 at *JXB* online).

## Discussion

### Chloroplast-dependent $^1\text{O}_2$ -mediated PCD in ACSC treated with RB

RB, which accumulates in chloroplasts of *Arabidopsis* cells, caused a surge in  $^1\text{O}_2$  production that activated transcriptional defence responses leading to the physiological induction of PCD. However, treatment with MV or IC, which do not enter chloroplasts, did not activate any statistical change in the transcriptional profile of ACSC. In spite of this, it could not be unambiguously concluded that the divergent subcellular location of the three  $^1\text{O}_2$  elicitors was the sole cause of this observation because the  $^1\text{O}_2$  quantum yields of the three elicitors are inherently different and, in particular, lower for MV and IC (Kovacs *et al.*, 2013). Therefore, this physiological defence response was investigated in more detail in both dark-grown and light-grown *Arabidopsis* cells in order to establish whether its activation was initiated by  $^1\text{O}_2$  production inside the chloroplasts or instead was independent of



**Fig. 6.** Overall picture of the hierarchical clustering analysis on the 485 transcripts with statistically differential expression and fold change  $|\text{Log}_2| > 1$  in ACSC after the treatment with 0.5  $\mu\text{M}$  RB. Data were clustered together with available expression microarray data from HL experiments and *Arabidopsis* mutants included in the GENEVESTIGATOR database: HL *hy5* vs LL *hy5*, *long hypocotyl5* transcription factor mutant of *A. thaliana* (Col-0) seedlings exposed to HL irradiance (1000  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) for 3h versus low light (LL) irradiance (100  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) (Kleine *et al.*, 2009); HL *npq1lut2* vs LL *npq1lut2*, double mutant of 6-week-old plants of *A. thaliana* (Col-0) lacking violaxanthin de-epoxidase and lycopene- $\epsilon$ -cyclase activities exposed to HL irradiance (1000  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) for 2h versus LL irradiance (25  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) (Alboresi *et al.*, 2011); HL CAT2HP1 vs LL CAT2HP1, 6-week-old catalase-deficient plants of *A. thaliana* (Col-0) exposed to HL irradiance (1600–1800  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) for 8h versus LL irradiance (100–140  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) (Vanderauwera *et al.*, 2005); RB-treated ACSC vs LL ACSC, this study; HL ACSC vs dark ACSC, *Arabidopsis* cell suspension cultures exposed to HL irradiance (1000  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) for 0.5h versus 1h dark conditions (González-Pérez *et al.*, 2011); HL-exposed plants vs LL plants; mature wild-type plants of *A. thaliana* (Col-0) exposed to HL irradiance (1400–1600  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) for 1h versus LL irradiance (40–70  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) (Rossel *et al.*, 2007); UV 310 nm vs 327 nm cut-off; seedlings of *A. thaliana* (Wassilewskija) exposed for 15 min at midday to UV-B from Philips TL 40W/12 UV fluorescent tubes unfiltered through a 3 mm quartz plate versus filtered through a 3 mm transmission WG327cut-off filter (Ulm *et al.*, 2004); *flu* vs *Ler*, the *flu* mutant of *A. thaliana* (*Ler*) exposed to light irradiance (90  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) for 2h after incubation in the dark for 8h versus dark conditions (Laloi *et al.*, 2007); *flu/over-tAPX* vs *over-tAPX*, the *flu* mutant of *A. thaliana* (*Ler*) overexpressing the thylakoid-bound ascorbate peroxidase (tAPX) exposed to light irradiance (90  $\mu\text{E m}^{-2} \text{s}^{-1}$ ) for 2h after incubation in the dark for 8h versus dark conditions (Laloi *et al.*, 2007). See the text for a detailed description of the main clusters A and B.

**Table 1.**  $^1\text{O}_2$  markers with statistically significant enrichment ( $P < 0.05$ ) in cluster A of the hierarchical clustering analysis of the 485 transcripts with fold change  $|\text{Log}_2| > 1$  in ACSC after the 30 min treatment with 0.5  $\mu\text{M}$  RB

| Element   | Locus ID               | Annotation   |
|-----------|------------------------|--|
| 248799_at | At5g47230              | ERF5, ethylene responsive element binding factor 5   |
| 266821_at | At2g44840              | ERF13, ethylene responsive element-binding factor 13   |
| 248448_at | At5g51190              | AP2 (Apeta2) domain-containing transcription factor, putative  |
| 262211_at | At1g74930              | Member of the dehydration response element-binding (DREB) subfamily A-5 of ERF/AP2 transcription factor family |
| 259992_at | At1g67970              | HSFA8, heat shock transcription factor A8  |
| 247707_at | At5g59450              | SCL11, Scarecrow-like transcription factor 11  |
| 252679_at | At3g44260              | CCR4-NOT transcription complex protein, putative   |
| 257511_at | At1g43000              | Zinc-binding family protein  |
| 252474_at | At3g46620              | Zinc finger (C3HC4-type RING finger) family protein  |
| 258436_at | At3g16720              | Zinc ion binding/protein binding   |
| 256306_at | At1g30370              | Lipase class 3 family protein  |
| 246600_at | At5g14930              | SAG101, senescence-associated gene 101; triacylglycerol lipase   |
| 245777_at | At1g73540              | NUDT21, Nudix hydrolase homologue 21   |
| 248793_at | At5g47240              | NUDT8, Nudix hydrolase homologue 8   |
| 252592_at | At3g45640              | MPK3, mitogen-activated protein (MAP) kinase 3   |
| 257751_at | At3g18690              | MKS1, MAP kinase substrate 1   |
| 255872_at | At2g30360              | CIPK11, Cystathionine $\beta$ -lyase-interacting protein kinase 11   |
| 251683_at | At3g57120 <sup>a</sup> | Protein kinase family protein  |
| 252470_at | At3g46930 <sup>a</sup> | Protein kinase family protein  |
| 263320_at | At2g47180              | GOLS1, galactinol synthase 1   |
| 247811_at | At5g58430              | EXO70B1, exocyst subunit EXO70 family protein B1   |
| 247240_at | At5g64660 <sup>a</sup> | U-box domain-containing protein  |
| 256999_at | At3g14200 <sup>a</sup> | DNAJ heat shock N-terminal domain-containing protein   |
| 246108_at | At5g28630              | Glycine-rich protein   |
| 266259_at | At2g27830              | Similar to pentatricopeptide (PPR) repeat-containing protein   |
| 246270_at | At4g36500 <sup>a</sup> | Similar to unknown protein (TAIR:At2g18210.1)  |
| 245840_at | At1g58420              | Similar to unknown protein (TAIR:At1g10140.1)  |

<sup>a</sup> Specific markers for  $^1\text{O}_2$  that are also up-regulated in ACSC after the treatment with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$ .

cellular location. It was determined that PCD was not activated by  $^1\text{O}_2$  produced by RB in dark-grown cells that had not developed chloroplasts; a result that was in full agreement with the recent study by Kim and co-workers (2012), who observed that the  $^1\text{O}_2$ -mediated cell death in seedlings of the leaf-variegated mutant *variegated2* was only induced in cells of the green leaf sector of this mutant containing fully developed chloroplasts, but not in the white sector where undifferentiated plastids were present. Chloroplasts are known both to be an important source of defence signalling molecules (i.e. ROS and precursors of defence hormones) and to have a central role in defence responses and the hypersensitive response (HR), a form of PCD localized at the site of pathogen attack that requires light for its development in many cases (Karpinski *et al.*, 2003; Coll *et al.*, 2011; Kacprzyk *et al.*, 2011). Likewise, the involvement of chloroplasts and ROS in PCD was recently demonstrated in ACSC (Doyle *et al.*, 2010), supporting the view that chloroplasts play a significant role in PCD regulation. In some studies carried out with *C. reinhardtii*, it has been proposed that  $^1\text{O}_2$  can escape from the chloroplasts and interact with an  $^1\text{O}_2$  sensor located in the cytoplasm or even reach the nucleus (Fischer *et al.*, 2007). However, on the basis of the present results with RB, which does not activate PCD in dark-grown cells, it is difficult to reconcile the results with a role for  $^1\text{O}_2$  itself as

a signalling molecule outside chloroplasts unless extremely, non-physiological HL conditions or very high concentrations of  $^1\text{O}_2$  elicitors come into play. Hence it is proposed that  $^1\text{O}_2$ -mediated PCD in ACSC is initiated only via developed chloroplasts.

Under the experimental conditions used here, 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  did not activate PCD in cells grown in dark or light conditions. Low PCD rates in ACSC subjected to  $\text{H}_2\text{O}_2$  were shown here; however, higher  $\text{H}_2\text{O}_2$  concentrations or longer incubation times than those described in this study were required to increase the PCD levels (Desikan *et al.*, 1998, 2001).

The functional enrichment analysis of the transcriptional profiling of light-grown ACSC also provided lines of evidence for the activation of PCD after the treatment with 0.5  $\mu\text{M}$  RB, but not with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$ . A direct comparison between the 30 min, 2-fold up-regulated transcripts in ACSC treated with 0.5  $\mu\text{M}$  RB and the 30 min early-induced, 2-fold up-regulated transcripts in the *flu* mutant (op den Camp *et al.*, 2003) showed that >40% of the up-regulated transcripts in ACSC (i.e. 138 of out 314) were included in the above list of up-regulated transcripts in the *flu* mutant. Likewise, the list of the 30 min, 2-fold, up-regulated transcripts in ACSC, treated with 0.5  $\mu\text{M}$  RB, contained 32 specific markers for  $^1\text{O}_2$  (Supplementary Table S6 at JXB online), seven specific markers for  $\text{H}_2\text{O}_2$ , and no specific markers for  $\text{O}_2^-$  when it



was compared with lists of specific markers for ROS provided by Gadjev and co-workers (2006). This indicated that  $^1\text{O}_2$  produced by RB in ACSC induced the expression of a list of transcripts that resembled the list observed in the *flu* mutant and confirmed that the transcriptional defence responses observed were mediated by  $^1\text{O}_2$  production inside chloroplasts. However, they did not seem to be associated with the EX1/EX2- and EDS1-dependent signalling pathway (see below). In addition, the analysis of the over-represented GO/biological process terms in *Arabidopsis* cells treated with 0.5  $\mu\text{M}$  RB showed that most of them corresponded to responses to pathogens, several biotic and abiotic stimuli, and PCD, which were also observed in the *flu* mutant (op den Camp *et al.*, 2003). Unexpectedly, the list of the 2-fold up-regulated transcripts in ACSC after the treatment with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  also exhibited some overlap (<30%) with the list of the 30 min early-induced, 2-fold up-regulated transcripts in the *flu* mutant and 19 specific markers for  $^1\text{O}_2$  (Supplementary Table S6), and only three specific markers for  $\text{H}_2\text{O}_2$  were identified. The fact that the treatment with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  inhibited the expression of several chloroplast transcripts ( $\text{Log}_2 < 1$ ) particularly associated with photosynthesis (Supplementary Table S3, locus identifier in a cyan background) suggested that  $\text{H}_2\text{O}_2$  clearly entered chloroplasts and activated transcriptional defence responses that had common characteristics with those described for the 0.5  $\mu\text{M}$  RB treatment and the *flu* mutant. This is confirmed in Supplementary Tables S7 and S8, where a significant number of over-represented GO/biological process terms after the treatments with 0.5  $\mu\text{M}$  RB and 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  appeared in both lists. However, a closer inspection of both lists disclosed that GO terms such as resistance response to pathogen (GO:0009814 and GO:009627), protein ubiquitination (GO:0016567), and PCD (GO:0012501 and GO:0006915) were only present in the treatment with 0.5  $\mu\text{M}$  RB. The list of the 30 min, up-regulated transcripts in ACSC after the treatment with 0.5  $\mu\text{M}$  RB contained key transcripts associated with the above GO terms that included: resistance (*R*) transcripts encoding TIR-NB-LRR-type resistance and CC-NB-LRR proteins (*At5g41750*; *At5g41740*; *At1g58807*; *At1g59124*; *At1g59620*), MITOGEN-ACTIVATED PROTEIN KINASES (*MPK3*, *At3g45640*; *MPKK4*, *At1g51660*), the ubiquitin ligase PLANT U-BOX17 (*PUB17*, *At1g29340*), a NUDIX HYDROLASE HOMOLOG (*NUDT7*, *At4g12720*), BCL-2-ASSOCIATED ATHANOGENE (*BAG3*, *At5g07220*), the lipase-like protein PHYTOALEXIN DEFICIENT 4 (*PAD4*, *At3g52430*), the ELICITOR PEPTIDE 2 PRECURSOR (*PROPEP2*, *At5g64890*), the transcription factor WRKY70 (*At3g56400*), and the ANKYRIN REPEAT-CONTAINING PROTEIN 2A (*AKR2A*; *At4g35450*); the underlined transcripts of which were also present in the early-induced, 2-fold up-regulated transcripts of the *flu* mutant. In contrast, the list of up-regulated transcripts in ACSC treated with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  contained only four transcripts associated with PCD and resistance response to pathogens (*BAG3*, *At5g07220*; *PROPEP2*, *At5g64890*; *AKR2A*, *At4g35450*; and *MYB30*, *At3g28910*), but none of them was included in the list of

the early-induced, 2-fold up-regulated transcripts in the *flu* mutant.

It has been proposed that ethylene (ET) and salicylic acid (SA) signalling pathways function additionally during the  $^1\text{O}_2$ -mediated PCD and that the oxylipins 12-oxophytodienoic acid (OPDA) and dinor-OPDA antagonize the cell death-inducing activity of jasmonic acid (JA) in the *flu* mutant (Danon *et al.*, 2005). In a search for key transcripts associated with the biosynthesis of the above enzymes in ACSC treated with 0.5  $\mu\text{M}$  RB, it was observed that 1-AMINOCYCLOPROPANE-1-CARBOXYLIC ACID (ACC) SYNTHASE 6 (*ACC6*, *At4g11280*) was present together with another eight ethylene-responsive element binding factors (ERFs) that included all those early-induced ERFs listed by Danon and co-workers (2005). In addition, transcripts encoding enzymes associated with the biosynthesis of OPDA in chloroplasts such as ALLENE OXIDE SYNTHASE (*AOS*, *At5g42650*), the conjugation of OPDA with glutathione such as GLUTATHIONE S-TRANSFERASE 5 (*GST5*, *At2g29450*), and the biosynthesis of JA in peroxisomes such as 3-oxo-2-[2'(Z)-pentenyl]-cyclopentane-1-octanoic (OPC-8:0) CoA LIGASE 1 (*OPCL1*, *At1g20510*) were also present. However, there was no evidence for the up-regulation of *EDS1* (*At3g48090*) known to be required for the biosynthetic activation of SA and the modulation of the  $^1\text{O}_2$ -mediated PCD in the *flu* mutant (Ochsenbein *et al.*, 2006). The essential regulation role of *EDS1* is mediated by its interaction with *PAD4*, and also by the product of the *SENESCENCE-ASSOCIATED GENE 101* (*SAG101*, *At5g14930*), two proteins that have defence regulatory functions beyond stabilizing *EDS1* in TIR-NB-LRR-type *R* gene-triggered resistance as demonstrated by silencing *EDS1* with a double-stranded DNA (dsRNA) interference construct (Feys *et al.*, 2005). The evidence that key transcripts associated with the biosynthesis pathway of ET and JA, together with *PAD4* and *SAG101*, are up-regulated in ACSC with 0.5  $\mu\text{M}$  RB suggests that they could be responsible for the induction of PCD through a signalling pathway that does not depend on *EDS1*. This agrees with the recent study by Ramel *et al.* (2013), who established that  $^1\text{O}_2$  activated cell death in the *chl1* mutant under HL stress, although no evidence for the activation of the EX1/EX2- and *EDS1*-dependent signalling pathway was found. In contrast, JA was described to have a key role in the  $^1\text{O}_2$ -mediated cell death, finding significant differences in the activation or repression of transcripts associated with the biosynthesis of this hormone under HL or acclimatory conditions. It is worth noting that an acclimatory response, but not PCD, was activated in ACSC subjected to a 30 min treatment with HL ( $1800 \mu\text{E m}^{-2} \text{s}^{-1}$ ) (González-Pérez *et al.*, 2011; Gutiérrez *et al.*, 2011) and that no evidence was found for the up-regulation of *EDS1*, *PAD4* and *SAG101* nor for transcripts encoding enzymes associated with the biosynthesis of JA in peroxisomes.

Similarly, a search for key early-induced transcripts associated with the biosynthesis and signalling cascade of ET, JA, and SA (listed in the study by Danon and co-workers, 2005) in ACSC treated with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  did not yield any match except for a transcript encoding the transcription factor ERF6 (*At4g17490*), supporting the view that PCD was

not activated either. Of interest is the observation that ACSC treated with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  exhibited several up-regulated transcripts encoding UDP-glycosyl transferases (UGTs) and GST, some of them known to be early-induced by  $\text{H}_2\text{O}_2$ , such as *UGT73B3*, *UGT73B5*, and *GST6* and with an important role in plant defence responses (Chen and Singh, 1999; Langlois-Meurinne *et al.*, 2005).

#### Correlation between the *flu* family mutants and ACSC treated with RB

The hierarchical clustering analysis of 485 co-regulated transcripts between ACSC treated with 0.5  $\mu\text{M}$  RB,  $^1\text{O}_2$ -producing mutants, and wild-type plants subjected to HL showed that ACSC treated with 0.5  $\mu\text{M}$  RB had higher Pearson's correlation with the *flu* family mutants than with the *npq1lut2* mutant. ACSC treated with 0.5  $\mu\text{M}$  RB had 27 specific markers for  $^1\text{O}_2$  in common with the *flu* mutant (Table 1), but only two out of the 27 were up-regulated in the *npq1lut2* mutant exposed to HL for 2 h (At5g47240 and At3g14200), where an acclimatory response was reported (Alboresi *et al.*, 2011). A manual inspection of the up-regulated transcripts (with  $\text{Log}_2$  ratios  $\geq 1$ ) in *chl1* (Ramel *et al.*, 2013) showed that four (At2g29450, At2g30360, At2g44840, and At3g44260) were listed among the 27 specific markers for  $^1\text{O}_2$  in cluster A when this mutant was subjected to HL for 2 d, but none when *chl1* was first acclimated to HL. Interestingly, the former up-regulated transcripts are involved in response to ET and JA stimuli. A transcriptional analysis of defence responses mediated by carotenoid oxidation products (i.e.  $\beta$ -cyclocitral) with origin in the reaction between  $^1\text{O}_2$  and the  $\beta$ -carotene molecules (housed in the PSII RC) also proved that  $^1\text{O}_2$  production by PSII RC induced an acclimatory rather than a PCD response (Ramel *et al.*, 2012a, b). A manual inspection of the up-regulated transcripts in the aforementioned study also indicated that only two out of the 27 specific markers for  $^1\text{O}_2$  in cluster A (At3g46620 and At3g14200) were up-regulated after the treatment with  $\beta$ -cyclocitral. In the study by Rosset and co-workers (2007), leaves were directly exposed to HL for 30 min. The list of up-regulated transcripts contained several specific markers for  $^1\text{O}_2$ , eight of which were included in cluster A (At1g74930, At2g30360, At2g35710, At2g44840, At2g47180, At3g44260, At3g46930, At5g47240, see Table 1 for their description). Despite the fact that this HL treatment induced a significant number of specific markers for  $^1\text{O}_2$ , the underlined transcripts of which were involved in response to ET and JA stimuli, the authors described the activation of a (systemic acquired) acclimatory response. The production of  $^1\text{O}_2$  and other ROS has also been demonstrated in plants exposed to high doses of UV, and their effects on the expression of transcripts encoding enzymes with an antioxidant role and with a UV-dependent expression have been recently investigated (Hideg *et al.*, 2013). However, the overlap between transcripts encoding several antioxidative enzymes in the *flu* mutant and the UV-induced transcripts was very limited. In fact, a single match with the transcript encoding GLUTAREDOXIN (*GRX480*, At1g28480)—also induced in other chemical treatments—was detected. A similar

conclusion was reached when the 14 UV-induced transcripts were compared with the 2-fold, up-regulated transcripts in ACSC treated with 0.5  $\mu\text{M}$  RB, as only *GRX480* was also up-regulated. Although the hierarchical clustering analysis shows certain correlation between experiments where  $^1\text{O}_2$  is produced, the genome-wide expression profiling and, consequently, the  $^1\text{O}_2$ -mediated defence responses (i.e. acclimation or PCD) vary significantly between experimental conditions, where the EX1/EX2- and EDS1-dependent signalling pathway is not activated or overlaps with other  $^1\text{O}_2$ -mediated signalling pathways belonging to a more complex signalling network.

## Conclusion

In brief, PCD has been activated in *Arabidopsis* cell cultures treated with submicromolar concentrations of RB, a potent, artificial  $^1\text{O}_2$  photosensitizer that accumulates inside chloroplasts. The fact that this  $^1\text{O}_2$ -mediated physiological response occurs in light-grown cultures, but not in dark-grown cultures, suggests that its activation requires functional chloroplasts. Additionally, a co-regulation analysis using the early-induced transcripts with a statistically significant fold change [ $\text{Log}_2$ ]  $> 1$  in ACSC treated with RB showed that this treatment had high correlation with the *flu* family mutants (although there was no evidence for the up-regulation of *EDS1*), but low correlation with other  $^1\text{O}_2$ -producing mutants or HL treatments in which an acclimatory response was activated instead of cell death.

## Supplementary data

Supplementary data are available at *JXB* online.

**Figure S1.** Volcano plots of the microarray experiments:

**Figure S2.** Validation of the microarray experiments in ACSC after the 30 min treatments with 0.5  $\mu\text{M}$  RB and 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$ .

**Table S1.** Transcripts and their corresponding primers selected for monitoring early ROS-mediated responses in ACSC under (photo)oxidative stress.

**Table S2.** Set of transcripts down- and up-regulated in ACSC after the 30 min treatment with 0.5  $\mu\text{M}$  RB (adjusted *P*-value  $< 0.05$ ).

**Table S3.** Set of transcripts down- and up-regulated in ACSC after the 30 min treatment with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  (adjusted *P*-value  $< 0.05$ ).

**Table S4.** Transcripts and their corresponding primers selected for validation of the microarray experiments in ACSC under (photo)oxidative stress.

**Table S5.** Fold changes ( $\text{Log}_2$ ) in the expression of selected transcripts responding to ROS production in ACSC after the 30 min treatment with RB, MV, and IC at 0.5  $\mu\text{M}$ , and  $\text{H}_2\text{O}_2$  at 500  $\mu\text{M}$ .

**Table S6.** Up-regulated specific markers for  $^1\text{O}_2$  in ACSC after several chemical treatments.

**Table S7.** Tree view of over-represented GO/biological process terms in *Arabidopsis* cell suspension culture after

the 30 min treatment with 0.5  $\mu\text{M}$  RB (adjusted  $P$ -value <0.05).

**Table S8.** Tree view of the over-represented GO/biological process terms in *Arabidopsis* cell suspension culture after the 30min treatment with 500  $\mu\text{M}$   $\text{H}_2\text{O}_2$  (adjusted  $P$ -value <0.05).

**Table S9.** Pearson's correlation between different experimental conditions and *Arabidopsis* mutants.

## Acknowledgements

This work is funded by the Junta de Castilla y León (ref: CSI002A10-2). CTD was funded by the Irish Research Council for Science, Engineering and Technology. The group is very grateful to the 'Institut de Biochimie & Physiologie Moléculaire des Plantes' (INIA), Montpellier, France for providing the cell culture. We also thank Mr José Javier Martín for technical assistance.

## References

- Alboresi A, Dall'Osto L, Aprile A, Carillo P, Roncaglia E, Cattivelli L, Bassi R.** 2011. Reactive oxygen species and transcript analysis upon excess light treatment in wild-type *Arabidopsis thaliana* vs a photosensitive mutant lacking zeaxanthin and lutein. *BMC Plant Biology* **11**, 62.
- Al-Shahrouf F, Arbiza L, Dopazo H, Huerta-Cepas J, Mínguez P, Montaner D, Dopazo J.** 2007. From genes to functional classes in the study of biological systems. *BMC Bioinformatics* **8**, 114.
- Al-Shahrouf F, Díaz-Uriarte R, Dopazo J.** 2004. FatiGO: a web tool for finding significant associations of Gene Ontology terms with groups of genes. *Bioinformatics* **20**, 578–580.
- Arellano JB, Li H, González-Pérez S, Gutiérrez J, Melø TB, Vacha F, Naqvi KR.** 2011. Trolox, a water-soluble analogue of  $\alpha$ -tocopherol, photoprotects the surface-exposed regions of the photosystem II reaction center *in vitro*. Is this physiologically relevant? *Biochemistry* **50**, 8291–8301.
- Axelos M, Curie C, Mazzolini L, Bardet C, Lescure B.** 1992. A protocol for transient gene expression in *Arabidopsis thaliana* protoplasts isolated from cell suspension cultures. *Plant Physiology and Biochemistry* **30**, 123–128.
- Benjamini Y, Hochberg Y.** 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**, 289–300.
- Benjamini Y, Yekutieli D.** 2001. The control of the false discovery rate in multiple testing under dependency. *Annals of Statistics* **29**, 1165–1188.
- Bolstad BM, Irizarry RA, Astrand M, Speed TP.** 2003. A comparison of normalization methods for high density oligonucleotide array data based on variance and bias. *Bioinformatics* **19**, 185–193.
- Brosche M, Schuler MA, Kalbina I, Connor L, Strid A.** 2002. Gene regulation by low level UV-B radiation: identification by DNA array analysis. *Photochemical and Photobiological Sciences* **1**, 656–664.
- Caldana C, Degenkolbe T, Cuadros-Inostroza A, Klie S, Sulpice R, Leisse A, Steinhauser D, Fernie AR, Willmitzer L, Hannah MA.** 2011. High-density kinetic analysis of the metabolomic and transcriptomic response of *Arabidopsis* to eight environmental conditions. *The Plant Journal* **67**, 869–884.
- Chang CC, Yang YT, Yang JC, Wu HD, Tsai T.** 2008. Absorption and emission spectral shifts of rose bengal associated with DMPC liposomes. *Dyes and Pigments* **79**, 170–175.
- Chen WQ, Singh KB.** 1999. The auxin, hydrogen peroxide and salicylic acid induced expression of the *Arabidopsis* *GST6* promoter is mediated in part by an ocs element. *Plant Journal* **19**, 667–677.
- Chomczynski P, Sacchi N.** 2006. The single-step method of RNA isolation by acid guanidinium thiocyanate–phenol–chloroform extraction: twenty-something years on. *Nature Protocols* **1**, 581–585.
- Coll NS, Epple P, Dangl JL.** 2011. Programmed cell death in the plant immune system. *Cell Death and Differentiation* **18**, 1247–1256.
- Cui XQ, Churchill GA.** 2003. Statistical tests for differential expression in cDNA microarray experiments. *Genome Biology* **4**, 210.
- Danon A, Miersch O, Felix G, op den Camp RG, Apel K.** 2005. Concurrent activation of cell death-regulating signaling pathways by singlet oxygen in *Arabidopsis thaliana*. *The Plant Journal* **41**, 68–80.
- Desikan R, Burnett EC, Hancock JT, Neill SJ.** 1998. Harpin and hydrogen peroxide induce the expression of a homologue of *gp91-phox* in *Arabidopsis thaliana* suspension cultures. *Journal of Experimental Botany* **49**, 1767–1771.
- Desikan R, Mackerness SAH, Hancock JT, Neill SJ.** 2001. Regulation of the *Arabidopsis* transcriptome by oxidative stress. *Plant Physiology* **127**, 159–172.
- Doyle SM, Diamond M, McCabe PF.** 2010. Chloroplast and reactive oxygen species involvement in apoptotic-like programmed cell death in *Arabidopsis* suspension cultures. *Journal of Experimental Botany* **61**, 473–482.
- Feyes BJ, Wiermer M, Bhat RA, Moisan LJ, Medina-Escobar N, Neu C, Cabral A, Parker JE.** 2005. *Arabidopsis* SENESCENCE-ASSOCIATED GENE101 stabilizes and signals within an ENHANCED DISEASE SUSCEPTIBILITY1 complex in plant innate immunity. *The Plant Cell* **17**, 2601–2613.
- Fischer BB, Hideg E, Krieger-Liszkay A.** 2013. Production, detection, and signaling of singlet oxygen in photosynthetic organisms. *Antioxidant and Redox Signaling* **18**, 2145–2162.
- Fischer BB, Krieger-Liszkay A, Hideg E, Snyrychova I, Wiesendanger M, Eggen RIL.** 2007. Role of singlet oxygen in chloroplast to nucleus retrograde signaling in *Chlamydomonas reinhardtii*. *FEBS Letters* **581**, 5555–5560.
- Gadjev I, Vanderauwera S, Gechev TS, Laloi C, Minkov IN, Shulaev V, Apel K, Inze D, Mittler R, van Breusegem F.** 2006. Transcriptomic footprints disclose specificity of reactive oxygen species signaling in *Arabidopsis*. *Plant Physiology* **141**, 436–445.
- Gálvez-Valdivieso G, Mullineaux PM.** 2010. The role of reactive oxygen species in signalling from chloroplasts to the nucleus. *Physiologia Plantarum* **138**, 430–439.
- González-Pérez S, Gutiérrez J, García-García F, Osuna D, Dopazo J, Lorenzo Ó, Revuelta JL, Arellano JB.** 2011. Early transcriptional defense responses in *Arabidopsis* cell suspension culture under high light conditions. *Plant Physiology* **156**, 1439–1456.
- Gutiérrez J, González-Pérez S, García-García F, Lorenzo Ó, Arellano JB.** 2011. Does singlet oxygen activate cell death in *Arabidopsis* cell suspension cultures? Analysis of the early transcriptional defense responses to high light stress. *Plant Signaling and Behavior* **6**, 1937–1942.
- Hideg E, Ayaydin F, Tandori J, Kalai T, Kovacs L.** 2011. Effects of singlet oxygen elicitors on plant leaves. Are these all caused by ROS? *European Biophysics Journal* **40**, 176–176.
- Hideg E, Jansen MAK, Strid Å.** 2013. UV-B exposure, ROS, and stress: inseparable companions or loosely linked associates? *Trends in Plant Science* **18**, 107–115.
- Hruz T, Laule O, Szabo G, Wessendorp F, Bleuler S, Oertle L, Widmayer P, Gruissem W, Zimmerman P.** 2008. Genevestigator V3: a reference expression database for the meta-analysis of transcriptions. *Advances in Bioinformatics* **2008**, 420747.
- Johnson RA, Wichern DW.** 1998. *Applied multivariate statistical analysis*, 2nd edn. Upper Saddle River, NJ: Prentice Hall.
- Jouanneau JP, Péaud-Lenoël C.** 1967. Growth and synthesis of proteins in cell suspensions of a kinetin dependent tobacco. *Physiologia Plantarum* **20**, 834–850.
- Kacprzyk J, Daly CT, McCabe PF.** 2011. The botanical dance of death: programmed cell death in plants. In: Kader JC, Delseny M, eds. *Advances in botanical research*. Amsterdam: Elsevier Ltd, 169–261.
- Karpinski S, Gabrys H, Mateo A, Karpinska B, Mullineaux PM.** 2003. Light perception in plant disease defence signalling. *Current Opinion in Plant Biology* **6**, 390–396.
- Kauss D, Bischof S, Steiner S, Apel K, Meskauskiene R.** 2012. FLU, a negative feedback regulator of tetrapyrrole biosynthesis, is physically linked to the final steps of the  $\text{Mg}^{2+}$ -branch of this pathway. *FEBS Letters* **586**, 211–216.
- Kim C, Apel K.** 2013. Singlet oxygen-mediated signalling in plants: moving from flu to wild type reveals an increasing complexity. *Photosynthesis Research* **116**, 455–464.



- Kim CH, Meskauskiene R, Apel K, Laloi C.** 2008. No single way to understand singlet oxygen signalling in plants. *EMBO Reports* **9**, 435–439.
- Kim C, Meskauskiene R, Zhang SR, Lee KP, Ashok ML, Blajicka K, Herrfurth C, Feussner I, Apel K.** 2012. Chloroplasts of *Arabidopsis* are the source and a primary target of a plant-specific programmed cell death signaling pathway. *The Plant Cell* **24**, 3026–3039.
- Kleine T, Kindgren P, Benedict C, Hendrickson L, Strand A.** 2007. Genome-wide gene expression analysis reveals a critical role for CRYPTOCHROME1 in the response of *Arabidopsis* to high irradiance. *Plant Physiology* **144**, 1391–1406.
- Kleine T, Voigt C, Leister D.** 2009. Plastid signalling to the nucleus: messengers still lost in the mists? *Trends in Genetics* **25**, 185–190.
- Kóvacs L, Ayaydin F, Kálai T, Tandori J, Kós, PB, Hideg E.** 2013. Assessing the applicability of singlet oxygen photosensitizers in leaf studies. *Photochemistry and Photobiology* (in press)
- Laloi C, Stachowiak M, Pers-Kamczyc E, Warzych E, Murgia I, Apel K.** 2007. Cross-talk between singlet oxygen- and hydrogen peroxide-dependent signaling of stress responses in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* **104**, 672–677.
- Langlois-Meurinne M, Gachon CMM, Saindrenan P.** 2005. Pathogen-responsive expression of glycosyltransferase genes *UGT73B3* and *UGT73B5* is necessary for resistance to *Pseudomonas syringae* pv *tomato* in *Arabidopsis*. *Plant Physiology* **139**, 1890–1901.
- Ledford HK, Chin BL, Niyogi KK.** 2007. Acclimation to singlet oxygen stress in *Chlamydomonas reinhardtii*. *Eukaryotic Cell* **6**, 919–930.
- Lee KP, Kim C, Landgraf F, Apel K.** 2007. EXECUTER1- and EXECUTER2-dependent transfer of stress-related signals from the plastid to the nucleus of *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* **104**, 10270–10275.
- Leisinger U, Rufenacht K, Fischer B, Pesaro M, Spengler A, Zehnder AJB, Eggen RIL.** 2001. The glutathione peroxidase homologous gene from *Chlamydomonas reinhardtii* is transcriptionally up-regulated by singlet oxygen. *Plant Molecular Biology* **46**, 395–408.
- Livak KJ, Schmittgen TD.** 2001. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. *Methods* **25**, 402–408.
- Lupinkova L, Komenda J.** 2004. Oxidative modifications of the photosystem II D1 protein by reactive oxygen species: from isolated protein to cyanobacterial cells. *Photochemistry and Photobiology* **79**, 152–162.
- McCabe PF, Leaver CJ.** 2000. Programmed cell death in cell cultures. *Plant Molecular Biology* **44**, 359–368.
- Medina I, Carbonell J, Pulido L, et al.** 2010. Babelomics: an integrative platform for the analysis of transcriptomics, proteomics and genomic data with advanced functional profiling. *Nucleic Acids Research* **38**, W210–W213.
- Meskauskiene R, Nater M, Goslings D, Kessler F, op den Camp RG, Apel K.** 2001. FLU: a negative regulator of chlorophyll biosynthesis in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* **98**, 12826–12831.
- Miyao M, Ikeuchi M, Yamamoto N, Ono T.** 1995. Specific degradation of the D1 protein of photosystem II by treatment with hydrogen peroxide in darkness: implications for the mechanism of degradation of the D1 protein under illumination. *Biochemistry* **34**, 10019–10026.
- Morrison H, Mohammad T, Kurukulasuriya R.** 1997. Photobiological properties of methylene violet. *Photochemistry and Photobiology* **66**, 245–252.
- Mullineaux PM, Baker NR.** 2010. Oxidative stress: antagonistic signaling for acclimation or cell death? *Plant Physiology* **154**, 521–525.
- Nakano R, Ishida H, Makino A, Mae T.** 2006. *In vivo* fragmentation of the large subunit of ribulose-1,5-bisphosphate carboxylase by reactive oxygen species in an intact leaf of cucumber under chilling-light conditions. *Plant and Cell Physiology* **47**, 270–276.
- Ochsenbein C, Przybyla D, Danon A, Landgraf F, Gobel C, Imboden A, Feussner I, Apel K.** 2006. The role of *EDS1* (enhanced disease susceptibility) during singlet oxygen-mediated stress responses of *Arabidopsis*. *The Plant Journal* **47**, 445–456.
- op den Camp RG, Przybyla D, Ochsenbein C, et al.** 2003. Rapid induction of distinct stress responses after the release of singlet oxygen in *Arabidopsis*. *The Plant Cell* **15**, 2320–2332.
- Petrov V, Vermeirssen V, De Clercq I, Van Breusegem F, Minkov I, Vandepoel K, Gechev TS.** 2012. Identification of *cis*-regulatory elements specific for different types of reactive oxygen species in *Arabidopsis thaliana*. *Gene* **499**, 52–60.
- Quinn GP, Keough MJ.** 2002. *Experimental design and data analysis for biologists*. Cambridge: Cambridge University Press.
- Ramel F, Birtic S, Cuine S, Triantaphylidès C, Ravanat J-L, Havaux M.** 2012a. Chemical quenching of singlet oxygen by carotenoids in plants. *Plant Physiology* **158**, 1267–1278.
- Ramel F, Birtic S, Ginies C, Soubigou-Taconnat L, Triantaphylides C, Havaux M.** 2012b. Carotenoid oxidation products are stress signals that mediate gene responses to singlet oxygen in plants. *Proceedings of the National Academy of Sciences, USA* **109**, 5535–5540.
- Ramel F, Ksas B, Akkari E, Mialoundama AS, Monnet F, Krieger-Liszky A, Ravanat JL, Mueller MJ, Bouvier F, Havaux M.** 2013. Light-induced acclimation of the *Arabidopsis chlorina1* mutant to singlet oxygen. *The Plant Cell* **25**, 1445–1462.
- Ramel F, Sulmon C, Serra AA, Gouesbet G, Couee I.** 2012. Xenobiotic sensing and signalling in higher plants. *Journal of Experimental Botany* **63**, 3999–4014.
- Reape TJ, Molony EM, McCabe PF.** 2008. Programmed cell death in plants: distinguishing between different modes. *Journal of Experimental Botany* **59**, 435–444.
- Roberts EL, Burguieres S, Warner IM.** 1998. Spectroscopic studies of indigo carmine dye in organized media. *Applied Spectroscopy* **52**, 1305–1313.
- Rossel JB, Wilson PB, Hussain D, Woo NS, Gordon MJ, Mewett OP, Howell KA, Whelan J, Kazan K, Pogson BJ.** 2007. Systemic and intracellular responses to photooxidative stress in *Arabidopsis*. *The Plant Cell* **19**, 4091–4110.
- Smyth GK.** 2004. Linear models and empirical Bayes methods for assessing differential expression in microarray experiments. *Statistical Applications in Genetics and Molecular Biology* **3**, article 3.
- Triantaphylidès C, Havaux M.** 2009. Singlet oxygen in plants: production, detoxification and signaling. *Trends in Plant Science* **14**, 219–228.
- Triantaphylidès C, Krischke M, Hoeberichts FA, Ksas B, Gresser G, Havaux M, Van Breusegem F, Mueller MJ.** 2008. Singlet oxygen is the major reactive oxygen species involved in photooxidative damage to plants. *Plant Physiology* **148**, 960–968.
- Ulm R, Baumann A, Oravec A, Mate Z, Adam E, Oakeley EJ, Schafer E, Nagy F.** 2004. Genome-wide analysis of gene expression reveals function of the bZIP transcription factor HY5 in the UV-B response of *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* **101**, 1397–1402.
- Vandenabeele S, Vanderauwera S, Vuylsteke M, Rombauts S, Langebartels C, Seidlitz HK, Zabeau M, Van Montagu M, Inze D, Van Breusegem F.** 2004. Catalase deficiency drastically affects gene expression induced by high light in *Arabidopsis thaliana*. *The Plant Journal* **39**, 45–58.
- Vanderauwera S, Zimmermann P, Rombauts S, Vandenabeele S, Langebartels C, Groussin W, Inze D, van Breusegem F.** 2005. Genome-wide analysis of hydrogen peroxide-regulated gene expression in *Arabidopsis* reveals a high light-induced transcriptional cluster involved in anthocyanin biosynthesis. *Plant Physiology* **139**, 806–821.
- Walters RG, Ibrahim DG, Horton P, Kruger NJ.** 2004. A mutant of *Arabidopsis* lacking the triose-phosphate/phosphate translocator reveals metabolic regulation of starch breakdown in the light. *Plant Physiology* **135**, 891–906.